


Biomimetic High-Valent Non-Heme Iron Oxidants for the *cis*-Dihydroxylation and Epoxidation of Olefins**

Jochen Bautz, Peter Comba,* Carlos Lopez de Laorden, Matthias Menzel, and Gopalan Rajaraman

High-valent iron oxidants are known to be the catalytically active species in heme enzymes, and the last decade has seen similar developments in non-heme-iron biomimetics and biochemistry.^[1,2] Extensive reactivity studies, structural, spectroscopic, and computational work have established {Fe^{III}-OOH} and {Fe^{IV}=O} as relevant intermediates in catalytic oxygenation processes.^[2,3] The species {Fe^V=O} has also been trapped, characterized, and found to be active in oxygen transfer, and discrete {Fe^V} species have been spectroscopically characterized.^[4,5] TauD, an α -ketoglutarate-dependent enzyme, is a well-characterized non-heme iron enzyme and has a high-spin ($S=2$) center in its active form.^[6] The aqua ion is the only other known high-spin ferryl complex.^[7,8] All biomimetic ferryl complexes characterized to date have intermediate-spin ($S=1$) electronic configuration. Herein we present experimental data of the iron-catalyzed oxidation of cyclooctene by H₂O₂ with the tetradentate bispidine ligand L¹ (Scheme 1a). The data indicate that unprecedented Fe^{IV}

With the pentadentate bispidine ligand L² (Scheme 1b) the {(L²)Fe^{II}}/H₂O₂ chemistry produced the low-spin {Fe^{III}OOH}, high-spin {Fe^{III}(O₂)}, and intermediate-spin ferryl {Fe^{IV}=O} complexes, which have been fully characterized.^[9,10] It was shown that there is a direct path from {(L²)Fe^{II}} to the {(L²)Fe^{IV}} ferryl complex, which was identified as the catalytically active species.^[10–12] In sharp contrast, in the corresponding L¹-based system, no Fe^{III} species has been identified, with the exception of an {Fe^{III}OOR} complex obtained by the oxidation of {(L¹)Fe^{II}} with *t*BuOOH; this complex was shown to be close to the spin-crossover limit.^[13] We have also been unable to trap and characterize the intermediate-spin ferryl complex. A DFT analysis indicated that this result is not unexpected, as the {(L¹)Fe^{II}} system is predicted to be directly oxidized by H₂O₂ to the novel {Fe^{IV}(OH)₂} complex with an intermediate-spin ($S=1$) electronic configuration, and which reacts through proton transfer to the corresponding and slightly more stable high-spin ($S=2$) ferryl oxidant.^[14]

Olefin oxidation experiments similar to those with the pentadentate ligand L² were carried out ((L¹)Fe^{II})/H₂O₂/cyclooctene = 1:10:1000, 298 K, aerobic or argon atmosphere, MeCN as solvent, TON_{max} = 10,^[15] reaction time 0.5 h, standard work-up procedure, and GC-MS analysis; for details see the Supporting Information). Product distribution and labeling studies (¹⁸O) indicate that, while the reactivities of the L¹- and L²-based catalysts are similar (TON_{total} = 3.8 and 5.0; Table 1), the reaction mechanisms must be different. With the L²-based system, diol products are only produced under argon and are stereochemically scrambled (*cis*- and



Scheme 1. Iron complexes of a) L¹ and b) L². X denotes coordinated solvent.

centers may be involved as the active oxidants, and these species are predicted by DFT calculations to be the novel intermediate-spin {Fe^{IV}(OH)₂} ($S=1$) and the high-spin {Fe^{IV}=O} ($S=2$) species, which have not been observed in model chemistry until now.^[7]

[*] Dr. J. Bautz, Prof. Dr. P. Comba, Dr. C. Lopez de Laorden, M. Menzel, Dr. G. Rajaraman
 Anorganisch-Chemisches Institut
 Universität Heidelberg, INF 270
 69120 Heidelberg (Germany)
 Fax: (+49) 6226-548-453
 E-mail: peter.comba@aci.uni-heidelberg.de

[**] Financial support by the German Science Foundation (DFG) and an Alexander von Humboldt fellowship to G.R. are gratefully acknowledged.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

Table 1: TON, product ratios, and ¹⁸O labeling studies for the iron-bispidine-catalyzed oxidation of cyclooctene with H₂O₂.^[a]

	{(L ¹)Fe}		{(L ²)Fe} ^[10]	
	O ₂	Ar	O ₂	Ar
diol, epoxide TON	1.6, 2.2	1.5, 1.0	0.0, 5.0	1.0, 1.0
epoxide % ¹⁸ O:				
H ₂ ¹⁸ O ₂ /H ₂ ¹⁸ O/ ¹⁸ O ₂	45/0/55	100/–/–	15/0/85	82/18/–
<i>cis</i> diol % ¹⁸ O from:				
H ₂ ¹⁸ O ₂ ^[b]	0/5/95	0/4/96	–	0/23/77
H ₂ ¹⁸ O ^[b]	92/8/0	93/7/0	–	0/30/0
<i>trans</i> diol % ¹⁸ O from:				
H ₂ ¹⁸ O ₂ ^[b]	–	–	–	0/56/44
H ₂ ¹⁸ O ^[b]	–	–	–	0/50/0

[a] TON_{max} = 10;^[15] error limit is 5%. [b] percentage of no/one/two ¹⁸O atoms in the diol product.


trans-diols). This result indicates a carbon-radical-based process.^[10] In the L¹-based system, there is a constant amount of pure *cis*-diol product, irrespective of whether oxygen is present or not. This finding suggests that there is a different reaction channel for the diol production with L¹, which does not involve carbon radical intermediates.

From the data of the {(L²)Fe}-based system in Table 1, it was concluded that, under an aerobic atmosphere, about 80% of the epoxide results from autoxidation, which arises from a carbon-based radical intermediate (5 vs 1 TON epoxide, in agreement with the labeling data), and this radical is also assumed to be responsible for the formation of diol (no diol in presence of O₂).^[10] For the {(L¹)Fe}-based system, there also is an increased amount of epoxide under aerobic conditions (TON 2.2 versus 1.0). This finding is in agreement with the fact that in air, 55% of the epoxide oxygen results from O₂ (¹⁸O labeling); the other 45% is from H₂O₂. The hypothesis for the L¹-based system is that the epoxide product is produced, as in the L²-based system, by direct attack of the ferryl oxygen at the double bond, and that this attack leads to a carbon-based radical intermediate. The fact that the diol production is not quenched by O₂ indicates that diol results from another reaction channel. This finding is supported by two observations, namely: a) there is only *cis*-diol product, and b) both oxygen atoms of the diol originate from the H₂O₂ oxidant. An appealing hypothesis is that the diol is produced by the new [(L¹)Fe(OH)₂]²⁺ oxidant.^[14] The experimentally determined mechanism of the L²-based system^[10] was confirmed by DFT calculations,^[16] and a similar set of calculations was therefore done with the L¹-based system to more thoroughly understand the new reaction mechanism, and specifically to validate the above hypotheses.

From the decay of the [(L¹)Fe^{II}(H₂O₂)(NCMe)₂]²⁺ precursor there are four possible high-valent iron complexes, [(L¹)Fe^V=O(OH)]²⁺, [(L¹)Fe^{IV}=O(OH)]⁺, [(L¹)Fe^{IV}(OH)₂]²⁺, and [(L¹)Fe^{IV}=O(OH₂)]²⁺; the latter is formed from the dihydroxo complex in a water-assisted proton-transfer reaction with an energy barrier of 34.7 kJ mol⁻¹.^[14] The potential oxidant [(L¹)Fe^V=O(OH)]²⁺ is not discussed in detail owing to unfavorable energetics.^[14,17]

Oxo-hydroxo-iron(IV) complexes, such as [(L¹)Fe^{IV}=O(OH)]⁺, are expected to perform both epoxidation and *cis*-dihydroxylation of olefins. This reactivity has been reported for the Fe-tpa-based system (tpa = tris-2-pyridylmethylamine).^[18,19] Epoxidation and dihydroxylation based on the [(L¹)Fe^{IV}=O(OH)]⁺ complex are exothermic but involve relatively high energy barriers (approx. 70 kJ mol⁻¹, see Supporting Information). Also, the direct formation of [(L¹)Fe^{IV}=O(OH)]⁺ has been predicted to be significantly less probable than that of [(L¹)Fe^{IV}(OH)₂]²⁺ and its tautomer [(L¹)Fe^{IV}=O(OH₂)]²⁺.^[14] Therefore, pathways arising from [(L¹)Fe^{IV}=O(OH)]⁺ are not considered as major reaction channels. Consequently, the hypothesis is that [(L¹)Fe^{IV}(OH)₂]²⁺, formed in a direct process from [(L¹)Fe^{II}(H₂O₂)]²⁺ by O–O bond homolysis, is the key intermediate in the oxidation processes (Scheme 2).

The key results of our DFT calculations related to the *cis*-diol production are illustrated in the energy diagram in Figure 1. [(L¹)Fe^{IV}(OH)₂]²⁺ has a triplet (S = 1) ground state



Scheme 2. Diol pathway (left) and epoxide pathway (right) for the *cis*-dihydroxylation and epoxidation of olefins.

with a margin of 18.2 kJ mol⁻¹ to the quintet (S = 2) state. The transition state of the concerted reaction to the *cis*-diol has a relative energy of 36.1 kJ mol⁻¹ on the triplet surface. The spin




Figure 1. Computed pathway for the iron-bispidine-catalyzed *cis*-dihydroxylation of ethylene (see also Scheme 2, left). The initial energy levels (left) refer to the triplet ground state (S = 1, 0.0) and quintet excited state (S = 2, 18.2 kJ mol⁻¹) of [(L¹)Fe^{IV}(OH)₂]²⁺. Spin densities and bond lengths are given in the Supporting Information.

density in the triplet transition state is distributed between the iron center (1.08), the two oxygen atoms (0.18, 0.19) and the partially oxidized olefinic carbon atoms (0.40, 0.27). The reaction is highly exothermic ($-272.3 \text{ kJ mol}^{-1}$) and the product has a quintet ground state. Therefore, the reaction involves a spin-crossover, which is expected to occur after the formation of the transition state (the optimized structure of the transition state on the quintet surface is given as Supporting Information).

The proton transfer from $[(L^1)\text{Fe}^{\text{IV}}(\text{OH})_2]^{2+}$ to $[(L^1)\text{Fe}^{\text{IV}}=\text{O}(\text{OH}_2)]^{2+}$ has an energy barrier of 34.7 kJ mol^{-1} and is exothermic ($-18.5 \text{ kJ mol}^{-1}$); this process also involves a spin-crossover from the $\{\text{Fe}^{\text{IV}}(\text{OH})_2\}$ ($S=1$) to the $\{\text{Fe}^{\text{IV}}=\text{O}(\text{OH}_2)\}$ ($S=2$) tautomer.^[14,20] Stepwise and concerted pathways have been considered for the epoxidation, and the results are presented in Figure 2. In the stepwise reaction on the quintet surface the energy barrier is 7.3 kJ mol^{-1} . Spin coupling between the two magnetic centers in the emerging Fe^{III} radical intermediate leads to several possible spin states, and a quintet state with high-spin Fe^{III} and spin-down on the radical center is found to be the ground state. We have also considered the possibility of a concerted transition state, which directly leads to the Fe^{II} -epoxide product. On the quintet surface we were not able to refine a true transition state. We have also tested for two-state reactivity by

computation of the reaction on the triplet-state surface. Both concerted and stepwise transition states are close-lying with energy barriers of 56.2 and 55.2 kJ mol^{-1} , respectively, but these barriers are so high in energy that they can be neglected.

The ring closure from the radical intermediate to form the Fe^{II} -epoxide product has an energy barrier of 16.1 kJ mol^{-1} on the quintet ($S=2$) surface, and the calculated energy barrier on the heptet ($S=3$) surface is prohibitively high ($233.1 \text{ kJ mol}^{-1}$). The final Fe^{II} -epoxide product also has a quintet ground state. In the final step the epoxide is cleaved, leading to the regeneration of the catalyst. This step has an energy barrier of 12.9 kJ mol^{-1} , and the overall reaction occurs in the quintet state. The reaction is exothermic in each step. From our previous DFT calculations with the iron complexes of the pentadentate ligand L^2 it emerges that the reaction of O_2 with the Fe^{III} radical intermediate has a lower barrier than the ring closure, and this is required for the formation of epoxide by an autoxidation process.^[16] A similar behavior is expected in this case.

The DFT calculations demonstrate that $[(L^1)\text{Fe}^{\text{IV}}(\text{OH})_2]^{2+}$ ($S=1$) may be responsible for both *cis*-dihydroxylation and epoxidation. Formation of *cis*-diol from intermediate-spin $[(L^1)\text{Fe}^{\text{IV}}(\text{OH})_2]^{2+}$ is relatively sluggish compared to the formation of epoxide from its high-spin tautomer (36.1 vs




Figure 2. Computed pathways (stepwise and concerted) for the iron-bispidine-catalyzed epoxidation of ethylene. The dashed line indicates a concerted transition state. See also Scheme 2, right. The initial energy levels (left) refer to the quintet ground state ($S=2$, 0.0) and triplet excited state ($S=1$, 11.5 kJ mol^{-1}) of $[(L^1)\text{Fe}^{\text{IV}}=\text{O}(\text{OH}_2)]^{2+}$. Spin densities and bond lengths are given in the Supporting Information.

23.4 kJ mol⁻¹, sum of energy barriers for each process). However, the generation of [(L¹)Fe^{IV}=O(OH₂)]²⁺ from [(L¹)Fe^{IV}(OH)₂]²⁺ by proton transfer also has a significant energy barrier (34.7 kJ mol⁻¹). Starting from [(L¹)Fe^{IV}(OH)₂]²⁺, the sum of the computed energy barriers reveals that *cis*-dihydroxylation is favored by 22 kJ mol⁻¹ over olefin epoxidation. Under anaerobic atmosphere this finding leads to the prediction of a diol/epoxide ratio of 1.6:1, in excellent agreement with the experimentally determined ratio of 1.5:1 (Table 1). Under aerobic conditions, the ratio cannot be directly compared because of the autoxidation process in the epoxidation path.

Our interpretation of the bispidine-iron-catalyzed olefin oxidation with H₂O₂ is at variance with the suggestion that the mechanism with the comparable tetradentate tpa- and bpmen-based catalysts (bpmen = *N,N'*-dimethyl-*N,N'*-bis(2-pyridylmethyl)-1,2-diaminoethane) involves a catalytically active {Fe^V=O} intermediate.^[21–25] Our proposal is based on {Fe^{IV}(OH)₂} (S = 1) and its tautomer {Fe^{IV}=O(OH₂)} (S = 2), which have not been considered and are derived from experiment and a DFT analysis. We suggest that the {Fe^V=O(OH)} pathway is less probable but cannot be excluded with our L¹-based system.^[14,17,26] There is no indication that our results and interpretation may be generalized, that is, it is quite possible that other systems follow different mechanistic pathways. An interesting recent observation, however, is that [(tpa)Fe^{IV}=O]²⁺, derived from [(tpa)Fe^{III}-OO*t*Bu]²⁺, is able to oxidize olefins.^[27]

Received: April 17, 2007

Revised: June 12, 2007

Published online: September 17, 2007

Keywords: dihydroxylation · epoxidation · iron · oxidation · reaction mechanisms

- [1] B. Meunier, S. P. de Visser, S. Shaik, *Chem. Rev.* **2004**, *104*, 3947.
- [2] M. Costas, M. P. Mehn, M. P. Jensen, L. Que, Jr., *Chem. Rev.* **2004**, *104*, 939.
- [3] "Metal-Oxo and Metal-Peroxo Species in Catalytic Oxidations": J.-J. Girerd, F. Banse, A. J. Simaan, *Struct. Bonding (Berlin)* **2000**, *97*, 145.
- [4] F. Tiago de Oliveira, A. Chanda, D. Banerjee, X. Shan, S. Mondal, L. Que, Jr., E. L. Bominaar, E. Münck, T. J. Collins, *Science* **2007**, *315*, 835.
- [5] J. F. Berry, E. Bill, E. Bothe, S. DeBeer George, B. Mienert, F. Neese, K. Wieghardt, *Science* **2006**, *312*, 1937.
- [6] J. C. Price, E. W. Barr, B. Tirupati, M. Bollinger, Jr., C. Krebs, *Biochemistry* **2003**, *42*, 7497.
- [7] O. Pestovsky, S. Stoian, E. L. Bominaar, X. Shan, E. Münck, L. J. Que, A. Bakac, *Angew. Chem.* **2005**, *117*, 7031; *Angew. Chem. Int. Ed.* **2005**, *44*, 6871.
- [8] Note that the fully characterized {Fe^{IV}=O} aqua ion (S = 2)^[7] is not considered to be a biomimetic ferryl complex.
- [9] M. R. Bukowski, P. Comba, C. Limberg, M. Merz, L. Que, Jr., T. Wüstuba, *Angew. Chem.* **2004**, *116*, 1303; *Angew. Chem. Int. Ed.* **2004**, *43*, 1283.
- [10] M. R. Bukowski, P. Comba, A. Lienke, C. Limberg, C. Lopez de Laorden, R. Mas-Ballesté, M. Merz, L. Que, Jr., *Angew. Chem.* **2006**, *118*, 3524; *Angew. Chem. Int. Ed.* **2006**, *45*, 3446.
- [11] J. Bautz, M. Bukowski, M. Kersch, A. Stubna, P. Comba, A. Lienke, E. Münck, L. Que, Jr., *Angew. Chem.* **2006**, *118*, 5810; *Chem. Int. Ed.* **2006**, *45*, 5681.
- [12] A. Anastasi, A. Lienke, P. Comba, H. Rohwer, J. E. McGrady, *Eur. J. Inorg. Chem.* **2007**, 65.
- [13] J. Bautz, P. Comba, L. Que, Jr., *Inorg. Chem.* **2006**, *45*, 7077.
- [14] P. Comba, G. Rajaraman, H. Rohwer, *Inorg. Chem.* **2007**, *46*, 3826.
- [15] Turnover number TON = 10 × (μmol product per μmol catalyst).
- [16] P. Comba, G. Rajaraman, unpublished results.
- [17] Formation of [(L¹)Fe^V=O(OH)] is thermodynamically and kinetically unfavorable with respect to the formation of [(L¹)Fe^{IV}(OH)₂] and [(L¹)Fe^{IV}=O(OH₂)], (all possible spin states and water-assisted as well as non-water-assisted pathways have been considered).^[13] Moreover, the first transition state of the reaction of [(L¹)Fe^V=O(OH)] with ethylene has a relative energy of 12 kJ mol⁻¹, which is slightly higher than the corresponding transition state of the [(L¹)Fe^{IV}=O(OH₂)]-based reaction (see Supporting Information).
- [18] The catalytically active species in the [Fe^{II}(tpa)(NCCH₃)₂]²⁺/H₂O₂ system leading to epoxide and diols has been proposed to be [(tpa)Fe^VO(OH)]²⁺.^[19]
- [19] R. Mas-Ballesté, M. Fujita, C. Hemmila, L. Que, Jr., *J. Mol. Catal. A* **2006**, *251*, 49.
- [20] This energy barrier has been calculated with Gaussian 03 (G03); there is a small difference to the basis sets used (TZVP vs. LACV3P**++), which is not expected to have a significant influence on the computed energy barriers.
- [21] A. Bassan, M. R. A. Blomberg, P. E. M. Siegbahn, L. Que, Jr., *J. Am. Chem. Soc.* **2002**, *124*, 11056.
- [22] A. Bassan, M. R. A. Blomberg, P. E. M. Siegbahn, L. Que, Jr., *Angew. Chem.* **2005**, *117*, 2999; *Angew. Chem. Int. Ed.* **2005**, *44*, 2939.
- [23] A. Bassan, T. Borowski, M. Lundberg, P. E. M. Siegbahn in *Concepts and Models in Bioinorganic Chemistry* (Ed.: H.-B. Kraatz, N. Metzler-Nolte), Wiley-VCH, Weinheim, **2006**, p. 63.
- [24] T. A. Jackson, L. Que, Jr. in *Concepts and Models in Bioinorganic Chemistry* (Ed.: H.-B. Kraatz, N. Metzler-Nolte), Wiley-VCH, Weinheim, **2006**, p. 259.
- [25] D. Quiñero, K. Morokuma, D. G. Musaev, R. Mas-Ballesté, L. Que, Jr., *J. Am. Chem. Soc.* **2005**, *127*, 6548.
- [26] Preliminary experiments on the dependence of water added to the catalyst solutions lead to significant differences in the total amount of products and product ratios; direct oxidation of the {(L¹)Fe^{II}} precursor with *t*BuOOH or iodosylbenzene diacetate only produces epoxide from cyclooctene. These preliminary experiments support our mechanistic proposal and show how the product selectivity can be tuned.
- [27] M. S. Seo, T. Kamachi, T. Kuouno, K. Murata, M. J. Park, K. Yoshizawa, W. Nam, *Angew. Chem.* **2007**, *119*, 2341; *Angew. Chem. Int. Ed.* **2007**, *46*, 2291.